


Mathematical questions and solutions Volume 6



Filesize: 1.19 MB

Reviews

*A high quality book as well as the font applied was fascinating to see. It generally fails to charge excessive. I am just effortlessly could possibly get a enjoyment of studying a composed book.
(Brant Dach)*

MATHEMATICAL QUESTIONS AND SOLUTIONS VOLUME 6



To get **Mathematical questions and solutions Volume 6** eBook, remember to access the link below and save the ebook or have access to additional information that are in conjunction with MATHEMATICAL QUESTIONS AND SOLUTIONS VOLUME 6 book.

RareBooksClub. Paperback. Book Condition: New. This item is printed on demand. Paperback. 32 pages. Dimensions: 9.7in. x 7.4in. x 0.1in. This historic book may have numerous typos and missing text. Purchasers can download a free scanned copy of the original book (without typos) from the publisher. Not indexed. Not illustrated. 1866 Excerpt: . . . of two of the diagonals of the same complete quadrilateral. Prove that if O_2, O_3 are their other points of intersection, then $OO_1(OO_j, OO_3$ are the tangents to the three conics at O . I. Solution by J. Dale; H. Tomlinson; and others. Let AA, BB, CC be the diagonals of the quadrilateral; A, B, C being collinear vertices; and O the common point. Draw the lines AOM , cutting BC in M ; COL , cutting AB in L ; and $LNMP$, cutting BC in N and AC in P ; also join ON, PB , and produce these lines to meet in O_3 . In the hexagon $AABBOO$, the lines $OA, BA; AB, O_3B; BA, O_2O$ intersect in the collinear points M, P, N ; therefore, by Pascals theorem, O_2 lies on the conic $AABBO$. In the hexagon $BBCCO_2$ the lines $OC, CB; CB, BO_2; BC, O_2O$; intersect in the collinear points L, P, M ; therefore O_2 lies on the conic $BBCCO$. In the pentagon $CCAAO$, the straight line OO_2 is drawn through O so that the lines $OC, CA; CA, AO; AC, OO_2$ intersect in the collinear points L, M, N ; therefore OO_2 is a tangent to the conic $CCAAO$. In the other two cases the proof is similar. Cor. 2. --Precisely the same work in tangential coordinates would lead to the reciprocal of this theorem, viz. : If $ABCD$ be a quadrilateral circumscribing a conic, and P, Q any two points on the conic, then a conic can be...



[Read Mathematical questions and solutions Volume 6 Online](#)



[Download PDF Mathematical questions and solutions Volume 6](#)

Related Books

**[PDF] Animalogy: Animal Analogies**

Follow the web link under to download and read "Animalogy: Animal Analogies" document.

[Save Book >](#)

**[PDF] The Whale Tells His Side of the Story Hey God, Ive Got Some Guy Named Jonah in My Stomach and I Think Im Gonna Throw Up**

Follow the web link under to download and read "The Whale Tells His Side of the Story Hey God, Ive Got Some Guy Named Jonah in My Stomach and I Think Im Gonna Throw Up" document.

[Save Book >](#)

**[PDF] Yearbook Volume 15**

Follow the web link under to download and read "Yearbook Volume 15" document.

[Save Book >](#)

**[PDF] Molly on the Shore, BFMS 1 Study score**

Follow the web link under to download and read "Molly on the Shore, BFMS 1 Study score" document.

[Save Book >](#)

**[PDF] The Secret Life of Trees DK READERS**

Follow the web link under to download and read "The Secret Life of Trees DK READERS" document.

[Save Book >](#)

**[PDF] The Mystery at Motown Carole Marsh Mysteries**

Follow the web link under to download and read "The Mystery at Motown Carole Marsh Mysteries" document.

[Save Book >](#)